

AD \_\_\_\_\_

USAARL REPORT NO. 70-1

MICROPUNCTURE STUDIES ON THE MECHANISM  
OF SULFATE EXCRETION BY THE RAT KIDNEY

By

S. W. Weinstein, MAJ, MC

JULY 1969

U. S. ARMY AEROMEDICAL RESEARCH LABORATORY  
Fort Rucker, Alabama



## DOCUMENT CONTROL DATA - R&amp;D

(Security classification of title, body of abstract and indexing annotation must be entered when the overall report is classified)

1. ORIGINATING ACTIVITY (Corporate author) U. S. Army Aeromedical Research Laboratory Fort Rucker, Alabama		2a. REPORT SECURITY CLASSIFICATION Unclassified	
		2b. GROUP	
3. REPORT TITLE MICROPUNCTURE STUDIES ON THE MECHANISM OF SULFATE EXCRETION BY THE RAT KIDNEY			
4. DESCRIPTIVE NOTES (Type of report and inclusive dates)			
5. AUTHOR(S) (Last name, first name, initial) Weinstein, Stephen W., MAJ, MC			
6. REPORT DATE July 1969		7a. TOTAL NO. OF PAGES 12	7b. NO. OF REFS 16
8a. CONTRACT OR GRANT NO.		9a. ORIGINATOR'S REPORT NUMBER(S) USAARL REPORT NO. 70 - 1	
b. PROJECT NO. 3AO62110A 819 c. Task 101 d.		(FY 70) 9b. OTHER REPORT NO(S) (Any other numbers that may be assigned this report)	
10. AVAILABILITY/LIMITATION NOTICES Distribution of this document is unlimited. Qualified requestors may obtain copies of this report from DDC.			
11. SUPPLEMENTARY NOTES		12. SPONSORING MILITARY ACTIVITY U. S. Army Medical R&D Command Washington, D. C. 20315	
13. ABSTRACT <p>A series of free flow micropuncture experiments were performed on rats undergoing sodium sulfate diuresis. End proximal tubular fluids and ureteral urines from the punctured kidney were collected. The data indicates that tubular fluid/plasma (TF/P) ratios for sulfate remained close to 1.0 and filtrate reabsorption was quantitatively normal during sulfate infusion in the proximal tubule. Final urine analysis suggested that all sulfate leaving the proximal convolution is excreted.</p> <p>The data is interpreted as evidence that sulfate is not handled by a <math>T_M</math> limited mechanism in the rat kidney. Rather it appears dependent upon filtration rate, proximal tubular reabsorptive rate and plasma concentration of the anion. A comparison of bicarbonate reabsorption during carbonic anhydrase inhibition to sulfate reabsorption in the rat nephron suggests greater proximal passive permeability to sulfate than bicarbonate and equally restricted distal nephron permeability.</p>			

14. KEY WORDS	LINK A		LINK B		LINK C	
	ROLE	WT	ROLE	WT	ROLE	WT
1. Renal Micropuncture						
2. Sulfate						
3. Nephron Permeability						
4. Ion Transport						

**INSTRUCTIONS**

**1. ORIGINATING ACTIVITY:** Enter the name and address of the contractor, subcontractor, grantee, Department of Defense activity or other organization (*corporate author*) issuing the report.

**2a. REPORT SECURITY CLASSIFICATION:** Enter the overall security classification of the report. Indicate whether "Restricted Data" is included. Marking is to be in accordance with appropriate security regulations.

**2b. GROUP:** Automatic downgrading is specified in DoD Directive 5200.10 and Armed Forces Industrial Manual. Enter the group number. Also, when applicable, show that optional markings have been used for Group 3 and Group 4 as authorized.

**3. REPORT TITLE:** Enter the complete report title in all capital letters. Titles in all cases should be unclassified. If a meaningful title cannot be selected without classification, show title classification in all capitals in parenthesis immediately following the title.

**4. DESCRIPTIVE NOTES:** If appropriate, enter the type of report, e.g., interim, progress, summary, annual, or final. Give the inclusive dates when a specific reporting period is covered.

**5. AUTHOR(S):** Enter the name(s) of author(s) as shown on or in the report. Enter last name, first name, middle initial. If military, show rank and branch of service. The name of the principal author is an absolute minimum requirement.

**6. REPORT DATE:** Enter the date of the report as day, month, year; or month, year. If more than one date appears on the report, use date of publication.

**7a. TOTAL NUMBER OF PAGES:** The total page count should follow normal pagination procedures, i.e., enter the number of pages containing information.

**7b. NUMBER OF REFERENCES:** Enter the total number of references cited in the report.

**8a. CONTRACT OR GRANT NUMBER:** If appropriate, enter the applicable number of the contract or grant under which the report was written.

**8b, 8c, & 8d. PROJECT NUMBER:** Enter the appropriate military department identification, such as project number, subproject number, system numbers, task number, etc.

**9a. ORIGINATOR'S REPORT NUMBER(S):** Enter the official report number by which the document will be identified and controlled by the originating activity. This number must be unique to this report.

**9b. OTHER REPORT NUMBER(S):** If the report has been assigned any other report numbers (*either by the originator or by the sponsor*), also enter this number(s).

**10. AVAILABILITY/LIMITATION NOTICES:** Enter any limitations on further dissemination of the report, other than those imposed by security classification, using standard statements such as:

(1) "Qualified requesters may obtain copies of this report from DDC."

(2) "Foreign announcement and dissemination of this report by DDC is not authorized."

(3) "U. S. Government agencies may obtain copies of this report directly from DDC. Other qualified DDC users shall request through \_\_\_\_\_."

(4) "U. S. military agencies may obtain copies of this report directly from DDC. Other qualified users shall request through \_\_\_\_\_."

(5) "All distribution of this report is controlled. Qualified DDC users shall request through \_\_\_\_\_."

If the report has been furnished to the Office of Technical Services, Department of Commerce, for sale to the public, indicate this fact and enter the price, if known.

**11. SUPPLEMENTARY NOTES:** Use for additional explanatory notes.

**12. SPONSORING MILITARY ACTIVITY:** Enter the name of the departmental project office or laboratory sponsoring (*paying for*) the research and development. Include address.

**13. ABSTRACT:** Enter an abstract giving a brief and factual summary of the document indicative of the report, even though it may also appear elsewhere in the body of the technical report. If additional space is required, a continuation sheet shall be attached.

It is highly desirable that the abstract of classified reports be unclassified. Each paragraph of the abstract shall end with an indication of the military security classification of the information in the paragraph, represented as (TS), (S), (C), or (U).

There is no limitation on the length of the abstract. However, the suggested length is from 150 to 225 words.

**14. KEY WORDS:** Key words are technically meaningful terms or short phrases that characterize a report and may be used as index entries for cataloging the report. Key words must be selected so that no security classification is required. Identifiers, such as equipment model designation, trade name, military project code name, geographic location, may be used as key words but will be followed by an indication of technical context. The assignment of links, rules, and weights is optional.

AD \_\_\_\_\_

USAARL REPORT NO. 70-1

MICROPUNCTURE STUDIES ON THE MECHANISM  
OF SULFATE EXCRETION BY THE RAT KIDNEY

By

S. W. Weinstein, MAJ, MC

JULY 1969

U. S. ARMY AEROMEDICAL RESEARCH LABORATORY  
Fort Rucker, Alabama

U. S. Army Medical Research and Development Command

Distribution Statement. Distribution of this document is unlimited.

## NOTICE

Qualified requesters may obtain copies from the Defense Documentation Center (DDC), Cameron Station, Alexandria, Virginia. Orders will be expedited if placed through the librarian or other person designated to request documents from DDC (formerly ASTIA).

### Change of Address

Organizations receiving reports from the U. S. Army Aeromedical Research Laboratory on automatic mailing lists should confirm correct address when corresponding about Laboratory reports.

### Disposition

Destroy this report when it is no longer needed. Do not return it to the originator.

### Distribution Statement

Distribution of this document is unlimited.

### Disclaimer


The findings in this report are not to be construed as an official Department of the Army position unless so designated by other authorized documents.

## ABSTRACT

A series of free flow micropuncture experiments were performed on rats undergoing sodium sulfate diuresis. End proximal tubular fluids and ureteral urines from the punctured kidney were collected. The data indicates that tubular fluid/plasma (TF/P) ratios for sulfate remained close to 1.0 and filtrate reabsorption was quantitatively normal during sulfate infusion in the proximal tubule. Final urine analysis suggested that all sulfate leaving the proximal convolution is excreted.

The data is interpreted as evidence that sulfate is not handled by a  $T_M$  limited mechanism in the rat kidney. Rather it appears dependent upon filtration rate, proximal tubular reabsorptive rate and plasma concentration of the anion. A comparison of bicarbonate reabsorption during carbonic anhydrase inhibition to sulfate reabsorption in the rat nephron suggests greater proximal passive permeability to sulfate than bicarbonate and equally restricted distal nephron permeability.

APPROVED:

  
ROBERT W. BAILEY  
COL, MSC  
Commanding

## TABLE OF CONTENTS

	Page
Introduction .....	1
Methods .....	2
Results .....	3
Discussion .....	9
References .....	11
Distribution List	
DD Form 1473	

---

## LIST OF ILLUSTRATIONS

<u>Figure</u>	Page
1. Correlation of Transit Time and Fractional Filtrate Reabsorption in the Proximal Convolute Tubule .....	5
2. Interrelation of Chloride and Inulin TF/P Ratios in the Proximal Convolute Tubule .....	7
3. Linear Correlation of Proximal Tubular Fractional Fluid Reabsorption and Overall Renal Sulfate Reabsorption .....	8

---

## TABLES

<u>Table</u>	Page
I. Proximal Tubular Data .....	4
II. Final Urine Data .....	6

# MICROPUNCTURE STUDIES ON THE MECHANISM OF SULFATE EXCRETION BY THE RAT KIDNEY

## INTRODUCTION

Sulfate excretion by the mammalian kidney appears governed by a classical tubular maximum ( $T_M$ ) limited mechanism (1). In clearance studies utilizing dogs (2) Lotspiech demonstrated a  $T_M$  for sulfate poised at a level approximating the plasma concentration of this anion. Schwartz et al (3) were unable to show a  $T_M$ . However, technical difficulties complicated their experiments. Pitts (4) has suggested that anion competition governs the sulfate  $T_M$ .

Chloride reabsorption appears governed primarily by the relative permeability of available anions and hydrogen ion in tubular fluid. Sodium ion provides the driving force (5). Bicarbonate is apparently less permeable than chloride except in the presence of abundant hydrogen ion (5) (6).

The requirement for maintenance of electrical neutrality in tubular fluid necessitates coupled movement of cations and anions in reabsorbate. Thus for each sodium ion reabsorbed, one anion must be accounted for either via a cation exchange or anion reabsorption. Since cation exchange and anion reabsorption are probably passive processes (5), relative permeabilities of the membrane to the various particles should govern their behavior.

With these thoughts in mind, the experiments included in this report were performed. Sulfate was infused intravenously to produce a diuresis in rats. End proximal tubular fractional reabsorption of filtrate and tubular fluid concentration of chloride were measured. The data indicates isotonic reabsorption of sulfate occurs in the proximal tubule under the conditions of these experiments. Distal nephron sulfate reabsorption is in contrast close to zero. The data thus indicate lack of a  $T_M$  for sulfate in the rat.



## METHODS

White Charles-River rats weighing 315-360 gms were anesthetized with Inactin 80 mg/kg. The trachea and both jugular veins were cannulated with polyethylene tubing. The left kidney was prepared for micropuncture as previously described (5) (7). The left ureter was cannulated with PE 10 polyethylene tubing. The surface tubules were transilluminated via a Zeiss Diascleral Cone adapted to a Zeiss Low Voltage Illuminator. No replacement fluid was given during the surgery.

One half hour before collections were commenced, an intravenous infusion of 0.5M  $\text{Na}_2\text{SO}_4$  was started at 0.05 ml/min. Carboxyl C14 Inulin  $10 \mu\text{c/ml}$  was contained in the infusion fluid. Just prior to collections the infusion rate was reduced to 0.025 ml/min.

End proximal tubular segments known to represent portions 50-60% of proximal length were selected (8) and free flow samples collected in sharpened single lumen pipettes utilizing Sudan Black stained light mineral oil blocks to prevent backflow contamination from distal portions of the tubule (15). A Leitz micromanipulator was utilized to perform the punctures.

Tail blood was collected before and after each puncture. Collection time and Lissamine Green transit time (8) to the puncture site were determined for each tubular fluid sample. Urines were collected from the ureteral cannula under oil in preweighed vials. Urine volume was then determined by weight. At the termination of the experiment a vena caval blood was collected.

Tubular fluids were measured for total volume in constant bore capillary tubing with an eyepiece micrometer. Aliquots of the fluid were analyzed for chloride via a modification of the electrotitrimetric technique of Ramsey, et al (9) (10). The remainder of the fluid was transferred to a Beckman Dioxane Cocktail (11) for C 14 counting in a Beckman Model 200 B Liquid Scintillation Counter.

Plasma and urine were counted in a similar manner. These were also analyzed for chloride via a Cotlove Chloridometer and sodium and potassium via flame photometry.

Tubular fluid to plasma ratios for inulin ( $\text{TF}/\text{P}_{\text{In}}$ ) and chloride ( $\text{TF}/\text{P}_{\text{Cl}}$ ) were calculated, as were inulin clearances ( $\text{C}_{\text{In}}$ ), sodium, potassium and chloride excretion rates, and individual nephron filtration rates.

Since techniques for sulfate measurement were not available, estimates of plasma and urine concentrations were calculated. Assumptions in these estimates were that plasma bicarbonate concentration was 25m M/L and other anions excluding sulfate and chloride approximated 10 meq/L. Since the sum of plasma cations equals the sum of plasma anions sulfate could be calculated from -

$$P_{Na^{+}} + P_{K^{+}} = P_{Cl^{-}} + P_{HCO_3^{-}} + P_{SO_4} + P_{other}.$$

For estimates of urinary sulfate excretion the sum of sodium and potassium excretion was utilized since the urine was almost entirely chloride free.

## RESULTS

Table I presents the tubular fluid data.  $TF/P_{Cl}$  values increase in these samples to the same extent as controls previously reported by this investigator (5). Nephron filtration rates and  $TF/P_{In}$  are within normal limits when compared to non-sulfate infused rats (5). Figure 1 presents the inulin data as a function of transit time to the pipette. Included are control data obtained during 0.1 ml/min. intravenous infusion of isotonic saline. No difference exists between the slopes for control and sulfate data. The half time for reabsorption is 9-10 seconds, a value in the range of the half time of 9.2 seconds determined in this Laboratory by the split droplet microperfusion technique utilizing isotonic saline (7).

Table II presents a summary of measurements made on the ureteral urine and plasma compared to control data collected during 0.1 ml/min IV isotonic saline.  $C_{In}$  was within normal limits compared to a previous control mean of 7.7 ml/min (12). Sodium and potassium excretion rates were 14.33  $\mu$ Eq/min and 2.74  $\mu$ Eq/min, respectively, compared to control values of 14.33  $\mu$ Eq/min and 1.86  $\mu$ Eq/min, values significantly elevated above control. Urine volume was .0321 ml/min and  $U/P_{In}$  was 60.

Figure 2 presents a plot of  $TF/P_{Cl}$  as a function of  $TF/P_{In}$ . Each point represents a single sample. An increment in  $TF/P_{In}$  correlates with a concomitant increase in  $TF/P_{Cl}$ . This suggests a direct proportion between fractional filtrate reabsorption and tubular fluid chloride concentration.

Figure 3 is a graph representing end proximal  $TF/P_{In}$  vs. calculated filtered over excreted sulfate. Each point represents tubular fluid and urine samples simultaneously collected. The line represents a 1:1 direct correlation.

Table 1

## Proximal Tubular Data

	Transit Time Seconds	TF/ P <sub>In</sub>	TF/ P <sub>C1</sub>	Nephron GFR m $\mu$ l/min
	8.6	1.26	1.11	34.4
	11.0	1.93	1.11	24.0
	7.5	1.34	1.08	35.2
	9.0	-	1.08	-
	9.5	1.43	1.18	19.5
	10.2	1.46	1.16	21.6
	8.0	1.78	1.15	38.5
	8.5	1.96	1.16	29.2
	10.5	2.89	1.18	37.5
	10.0	2.36	1.19	35.2
	13.0	3.99	1.21	56.4
	15.0	4.96	1.19	46.7
	16.0	4.55	1.20	40.4
	16.0	3.65	1.17	19.5
Mean $\pm$ SEM	10.9 $\pm$ 0.8	2.58 $\pm$ 0.34	1.16 $\pm$ 0.01	33.7 $\pm$ 2.9
Control $\pm$ SEM (5) (12)	8.8 $\pm$ 0.5 (18)	1.82 $\pm$ 0.14 (23)	1.15 $\pm$ 0.02 (8)	31.3 $\pm$ 3.4 (17)

$\frac{P}{TF}$   
Inulin




Figure 1

Table II

## Final Urine Data

	$C_{In}$ ml/min	Urine Volume ml/min	$U/P_{In}$	$U_{Na} V$ $\mu Eq/min$	$U_K V$ $\mu Eq/min$
9	10.4	.0169	97	8.72	3.28
	11.1	.0145	120	8.16	3.15
	9.2	.0130	111	7.83	2.82
	10.3	.0129	126	6.49	2.80
	10.8	.0523	35	23.71	2.92
	14.5	.0461	53	22.00	2.94
	7.6	.0344	37	16.69	3.04
	8.2	.0334	42	15.10	2.85
	8.1	.0358	38	17.50	2.86
	8.3	.0430	35	17.89	2.65
	9.5	.0409	42	16.32	2.38
	8.1	.0381	38	14.48	2.29
	7.8	.0384	36	14.36	2.42
	5.0	.0306	29	11.38	2.01
Mean $\pm$ SEM	$9.2 \pm 0.57$	$.0321 \pm .003$	$60 \pm 9.3$	$14.33 \pm 1.36$	$2.74 \pm 0.09$
Control Mean $\pm$ SEM (12)	$7.7 \pm 0.42$	$.0107 \pm .003$	$82.6 \pm 11.6$	$1.28 \pm 0.35$	$1.86 \pm 0.23$




Figure 2




Figure 3

The data closely approximate that line.

## DISCUSSION

These results provide evidence suggesting that sulfate reabsorption in the proximal tubule of the rat kidney is not  $T_M$  limited. The chloride ratios from the end proximal samples continue to rise quantitatively to the same degree as during control conditions. If proximal sulfate reabsorption were significantly limited, its concentration might increase to exceed plasma values as filtered water was reabsorbed. Since bicarbonate reabsorption must continue in the presence of normal hydrogen secretion, the rise in sulfate would replace the anion deficit produced by dissipation of bicarbonate. Chloride would either remain close to its plasma concentration or possibly fall below that level. Thus it is concluded that sulfate concentration did not rise significantly above plasma levels. The correlation between  $TF/P_{In}$  and  $TF/P_{Cl}$  lends further support to this conclusion as one would anticipate a fall in chloride or minimal rise with increased filtrate reabsorption if sulfate concentration were increasing in tubular fluid.

Another mechanism for manifesting decreased proximal tubular sulfate reabsorption would be a decrease in the fractional and/or absolute reabsorption of filtered volume. Nephron filtration rate and proximal fractional reabsorption were within normal limits during sulfate infusion. Thus volume reabsorption was not reduced. Since  $TF/P$  for sulfate approximates 1.0 and volume reabsorption is normal in the proximal tubule, sulfate must be readily reabsorbed in this segment at a concentration close to that in plasma.

The measurements in ureteral urine from the experimental kidney support and extend the micropuncture data. As indicated in Figure 3, the fraction of sulfate reabsorbed in final urine closely approximates the fraction of filtrate reabsorbed in the end proximal tubular segments. Since the sulfate  $TF/P$  approaches 1.0 in these segments the fraction of filtered sulfate and water reabsorbed proximally must be equal. Any further sulfate reabsorption along more distal portions of the nephron should be reflected in the ureteral urine. The measured fraction of filtered sulfate reabsorbed in ureteral urine and in end proximal tubular segments are nearly equal. Thus little or no net sulfate reabsorption seems to occur in the distal nephron in these experiments.

This would appear in conflict with the data of Lotspiech (2) who clearly demonstrated a  $T_M$  in dogs for sulfate at plasma sulfate concentrations well



below those noted in the present experiments. Schwartz, et al (3) were unable to demonstrate a sulfate  $T_M$  in dogs. They observed a continued reabsorption of sulfate over a wide range of plasma concentration of this anion. Since they used a single intravenous injection of sodium sulfate, constant plasma levels of the anion were not achieved. This rendered a more quantitative analysis of the data difficult.

These micropuncture results are not consistent with a  $T_M$  for sulfate in the rat. They suggest that sulfate reabsorption is determined by filtration rate, plasma sulfate concentration, and proximal tubular reabsorptive capacity. The remainder of the nephron appears almost if not entirely impermeable to sulfate since little or no net reabsorption of this anion occurs beyond the proximal convolution.

Comparing these results to measurements of bicarbonate reabsorption during carbonic anhydrase inhibition is of interest. With reduced hydrogen ion secretion bicarbonate concentration rises above plasma in the proximal tubule (13) and TF/P for chloride falls to values significantly below 1.0 (5). This coincides with fractional and absolute proximal filtrate reabsorption not statistically different from control values (5) (16). Thus proximal bicarbonate reabsorption is depressed by carbonic anhydrase inhibition. In contrast sulfate reabsorption in the proximal tubule is unrestricted. This suggests the proximal tubule is more permeable to sulfate than bicarbonate. Final urine measurements for bicarbonate excretion during carbonic anhydrase inhibition suggest that no net bicarbonate reabsorption occurs beyond the proximal tubule. Thus the distal nephron appears equally impermeable to both sulfate and bicarbonate.

In summary, the data presented provide evidence that sulfate is not handled by a  $T_M$  limited mechanism in the rat kidney. Rather, it appears reabsorbed isototically in the proximal tubule over wide ranges of plasma sulfate concentration. Distal nephron sulfate reabsorption seems markedly limited such that most, if not all, of this anion leaving the proximal tubule is excreted in the final urine. A comparison of bicarbonate and sulfate reabsorption suggests greater passive permeability to sulfate than to bicarbonate in the proximal tubule. Distal permeability appears equally limited for both anions.

## REFERENCES

1. Smith, H. W. The Kidney. Structure and Function in Health and Disease. Oxford Univ. Press, N. Y., 1951, pgs 121-126.
2. Lotspiech, W. D. Renal Tubular Reabsorption of Inorganic Sulfate in the Normal Dog. Am. J. Physiol. 151:311, 1947.
3. Schwartz, B. M., P. K. Smith, and A. W. Winkler. Renal Excretion of Sulfate. Am. J. Physiol. 137: 658, 1942.
4. Pitts, R. F. Physiology of the Kidney and Body Fluids. Year Book Medical Publ., Chicago, 1963, pgs 77-78.
5. Weinstein, S. W. Micropuncture Studies of the Effects of Acetazolamide on Nephron Function in the Rat. Am. J. Physiol. 214:222, 1968.
6. Gottschalk, C. W. Renal Tubular Function: Lessons from Micropuncture. Harvey Lecture Ser. 58:99, 1963.
7. Weinstein, S. W. Effects of Cyanide and 2 Deoxyglucose on Proximal Tubular Function in the Rat Kidney. USAARL Report No. 69-17.
8. Steinhausen, M. Eine Methode zur Differenzierung proximaler und distaler Tubuli der Nierenrinde von Ratten in vivo und ihre Anwendung zur Bestimmung tubulärer Strömungsgeschwindigkeiten. Pflügers Arch ges Physiol. 277:23, 1963.
9. Ramsay, J., R. H. J. Brown and P. C. Croghan. Electrometric Titration of Cl in Small Volumes. J. Exp. Biol. 32:822, 1955.
10. Windhager, E. E. and G. Giebisch. Micropuncture Study of Renal Tubular Transfer of Sodium Chloride in the Rat. Am. J. Physiol. 200:581, 1961.
11. Beckman Instruction Manual. Model 200-B Liquid Scintillation Counter.

12. Weinstein, S. W. and R. M. Klose. Micropuncture Studies on Metabolism and Sodium Transport in the Mammalian Nephron. Am. J. Physiol. 217:1969.
13. Clapp, J. R., J. F. Watson, and R. W. Berliner. Effect of Carbonic Anhydrase Inhibition on Proximal Tubular Bicarbonate Reabsorption. Am. J. Physiol. 205:273, 1963.
14. Bernstein, B. A. and J. R. Clapp. Micropuncture Study of Bicarbonate Reabsorption by the Dog Nephron. Am. J. Physiol. 214:251, 1968.
15. Windhager, E. E. Micropuncture Techniques and Nephron Function. Butterworths, London, 1968. Chapters 3 & 4.
16. Dirks, J. H., W. J. Cirkse, and R. W. Berliner. Micro-puncture Study of the Effect of Various Diuretics on Sodium Reabsorption by the Proximal Tubules of the Dog. J. Clin. Invest. 45:1875, 1966.